

Published 2006

Copies of this map may be obtained from:
Information Sales Office Geological Survey of Canada
Alberta Geological Survey 1-888-252-4301 (toll free in Canada and USA)
Telephone: (780) 422-3767 Available at www.nrcan.gc.ca/gscbookstore
Website: www.agc.gov.ab.ca

AGS Map 360 GSC Open File 5183


Surficial Geology of the Beatty Lake Area (NTS 84M/NE)

Geology by: R.C. Paulen, A. Plouffe and I.R. Smith

Scale 1:100 000
2 0 2 4 6 8 10 kilometres
1 0 1 2 3 4 5 6 miles

Projection: Universal Transverse Mercator
Datum: North American Datum, 1983

Canada



Index to adjacent sheets	
GSC OF 5070	AGS Map 360
AGS Map 395	GSC OF 5183
84M	
GSC OF 5237	AGS Map 361
AGS Map 396	GSC OF 5184

OPEN FILE DOSSIER PUBLIC 5183	Open files are products that have not gone through the GSC formal publication process.
GEOLOGICAL SURVEY OF CANADA COMMISSION GÉOLOGIQUE DU CANADA 2006	Les dossiers publics sont des documents qui n'ont pas encore suivi le processus officiel de publication de la CGC.

Recommended reference format:
Paulen, R.C., Plouffe, A. and Smith, I.R. (2006): Surficial geology of the Beatty Lake area, Alberta (NTS 84M/NE); Alberta Energy and Utilities Board, EUB/AGS Map 360 and Geological Survey of Canada, Open File 5183, scale 1:100 000.


This is a common map legend for the surficial geology of northern Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

UNIT


UNIT NAME

DESCRIPTION AND GENESIS	
QUATERNARY HOCLOCENE	
A	ANTHROPOGENIC MATERIALS: Culturally made or modified geological materials such that their physical properties (e.g., structure, cohesion, compaction) have been drastically altered.
O	ORGANIC DEPOSITS: Undifferentiated peat (woody to fibrous muck) occurring in undifferentiated wetlands; commonly underlain by fine-grained, poorly drained glaciolacustrine deposits; includes marshes, swamps, bogs and fens. Bog peat: Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is dominated by sphagnum mosses, heath shrubs and short, stunted trees. Fen peat: Occurs in a peatland with water table at surface and slow internal drainage; peatland surface is dominated by sedges, with grasses and reeds near local pools, and is sparsely treed.
OB	COLLUVIAL DEPOSITS: Materials that have reached their present position as a result of direct, gravity-induced movement; commonly occurs as slope and slump deposits confined to valley slopes and floors; includes pre-existing bedrock, till, glaciolacustrine, glaciofluvial and eolian sediments, generally poorly sorted.
OF	FLUVIAL DEPOSITS: Sediments transported and deposited by streams and rivers; synonymous with alluvium. Includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g., postglacial floodplains, terraces, fans and deltas).
C	LACUSTRINE DEPOSITS: Sediments deposited in and adjacent to recent and modern lakes; offshore sand, silt and clay, minor organic deposits; littoral (nearshore) beaches and bars; sand, silt and minor gravel.
F	ELIAN DEPOSITS: Wind-deposited sediments; well-sorted, medium to fine-grained sand and minor silt (loess); generally massive to locally cross bedded or ripple laminated; includes both active and vegetated deposits.
L	GLACIOLACUSTRINE DEPOSITS: Primarily fine-grained, distal sediments deposited in or along the margins of glacial lakes, including sediments released by the melting of floating ice. Includes laminated (rhythmically bedded) to massive fine sand, silt and clay, and may contain ice-rafted stones.
E	LITTORAL AND NEARSHORE SEDIMENTS: Massive to stratified, well-sorted silty sand, pebbly sand and minor gravel; occurs as beaches, bars, spits and forest deltaic deposits deposited during regression and lowering of glacial lakes.
LG	GLACIOFLUVIAL DEPOSITS: Sediments deposited by glacial meltwater streams as suberial or subaqueous outwash. Includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Includes meltwater channels, kettle holes, depressions and/or ice-contact ridges.
LGL	ICE-CONTACT SEDIMENTS: Sediments deposited by glacial meltwater streams in direct contact with glacial ice, or from (kettle) terraces (e.g., crevase ridges) glacial ice. Includes massive to stratified, poor to moderately sorted, coarse sediments (predominantly pebble gravel and coarse sand, locally till) and may show evidence of ice melting (slumped structures).
FG	MORAINE: Nonsorted till (til) deposited directly by glacial ice consisting of a mixture of clay, silt, sand and minor pebbles, cobbles and boulders. Locally, this unit may contain blocks of bedrock, pre-existing stratified sediment and till, and lenses of glaciolacustrine and/or glaciocluvial sediment.
FGL	Stagnant ice moraine: Material resulting from the collapse and lateral movement of glacial ice margin; sediment is mainly diamict, but locally includes stratified sediments of glaciolacustrine or glaciocluvial origin. Characterized by low to high relief hummocky topography.
MT	Ice-thrust terrace: Terrain resulting from glacio-tectonic transport of originally subglacial material deposited by the greater or lesser extent; may include syn genetic till, as well as masses of pre-existing till, slumped drift and bedrock. Characterized by high to moderate relief and features include hill-pans and glacio-tectonic moraine ridges.
MF	Fluted moraine: Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smoothed hills, all landforms parallel the local ice flow direction; includes flutes, drumlins and drumlinoths.
FP	PREGLACIAL FLUVIAL DEPOSITS: Sediments transported and deposited by streams and rivers prior to glaciation. Includes sand and gravel deposits occurring in paleovalleys (i.e., preglacial floodplains, terraces, fans and deltas); ranging from middle Wisconsin to late Tertiary.
RT	UNCONSOLIDATED FLUVIAL GRAVELS: Predominately well-sorted, quartzite and chert gravel and cobbles; Cordilleran source, Paleogene (Tertiary) to early Quaternary.
R	BEDROCK: Undifferentiated; may include clastic sedimentary rock, shale, coal, carbonate and crystalline (Shield), kimberlite and/or coal.

SYMBOL LEGEND



BASEMAP LEGEND



UNIT NOTATION

Example: GLACIOLACUSTRINE plain

Textural modifier
s = sandy
G = gravel
L = loamy
C = clay
a = sand-silt-clay
Genetic modifier
p = pebble
g = gravel
s = sand
t = till
c = clay
a = sand-silt-clay

TEXTURAL MODIFIER

Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture; i.e., sc for sandy clay

p = pebble
g = gravel
s = sand
t = till
c = clay
a = sand-silt-clay

GENETIC & GEOMORPHIC MODIFIERS

c = crevass fill
d = doughnut rings circular hummocks with a central depression, plateau mounds and brain-like pattern ridges,
e = eroded
f = fan
g = gullied
h = hummock
k = collapse
m = meander
p = plain
r = ridged
s = slumped
t = terrace
u = undulating
v = veneer
w = washboard
z = dissected
l = delta
ice-contact ridges, ice-squeeze deposits and linear forms deposited by meltwater in stagnant ice
low to moderate relief
planar surface eroded by glacial meltwater, often capped by a boulder lag deposit and/or thin deposit of sand and gravel
gently sloping fan-shaped mass of detrital debris
slopes dissected by modern ravines created by intermittent runoff
assemblage of approximately equidimensional hills and hollows; moderate to high relief (commonly greater than 2 m)
depression, including kettles, pitted morphology, thermokarst depressions, karst sinkholes
sinuous curves, loops and oxbows produced as meltwater and modern streams shift their channels over time
deposit greater than 2 m thick; commonly masks geomorphic pattern of underlying deposits; flat to gently rolling topography (commonly less than 2 m relief)
one or more parallel or subparallel, convex, linear morphological elements with a length-to-width ratio greater than 2: low to high relief
landslide blocks, slope failure debris
terrace bench cut by either meltwater or wave action; antiplanation terrace, kame terrace
low-relief rolling terrain; swell and swale topography
thin mantle of unconsolidated material too thin to mask the minor irregularities of the surface of the underlying material; it ranges in thickness from 10 cm to 1 metre and may be discontinuous
low-relief transverse moraine ridges, usually formed from basal ice shearing
channelled or dissected by glacial meltwater flow; dissected terrain by Holocene fluvial activity
lake delta; ice-contact delta

COMPLEX

Where two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the combination is given in a two or three position designation set off by slashes denoting arbitrary percentage limits. For example,

'Mp/LGv' means the area is underlain by approximately 60% morainal plain and up to 40% glaciolacustrine veneer.

'MvLGl/Gp' means at least 60% of the area is underlain by morainal veneer, with up to 40% glaciolacustrine veneer and less than 15% glaciocluvial plain.

'LGp/M' means more than 60% of the area is underlain by a glaciolacustrine plain, with less than 15% moraine.

Stratigraphic Sequence

Where materials of different origins or textures are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:

'SLGv | Mp' Thin sandy glaciolacustrine sediment deposited on morainal plain

Transitional Association

Locally, two or more terrain units are juxtaposed by reason of related origin, temporal sequence or ambiguous geometric distinction. In the last case, both components may be present. Such situations are identified by a compound designation marked by a hyphen. Examples: 'FGz-LGz' indicating ice-contact delta distinguishing from glaciolacustrine delta, or 'FGk-MSh' indicating ice-contact kette and kettle topography that blends with hummocky stagnant ice moraine.

Morphologic Overprint

Where a sequence of geomorphic processes has produced a multi-aspect or compound terrain fabric, the geomorphic modifier suffixes are appended in the inferred order of superposition. Mp' means a plain of till has been moulded into ridge forms and finally dissected by modern streams. FGphr' means a glaciocluvial plain has been discontinuously covered by ice-contact hummocks and ridges.

Acknowledgements:

Surficial mapping was completed in 2004 as an Alberta Geological Survey (AGS) and Geological Survey of Canada (GSC) collaborative contribution to NRD Project 4450 and the Quaternary mapping initiative under the Alberta Mineral Strategy. Ryan Peterson assisted with fieldwork and database compilation. Paramount Resources Ltd. is gratefully acknowledged for providing logistical support at the remote Paramount Bistcho gas plant and camp. Field visits by Laurence Andriashuk and Rick Richardson provided insight into the permafrost development and aggregate inventory of the region. Digital cartography and GIS were completed by Monica Price and GiMo Solutions Ltd. Base data provided by Spatial Data Warehouse Ltd.

References:


- Edwards, W.A.D., Budney, H.D., Bereznik, T. and Butkovic, L. (2004): Sand and gravel deposits with aggregate potential, Bistcho Lake, Alberta (NTS 84M/6). Alberta Energy and Utilities Board, EUB/AGS Map 310, scale 1:250 000.
- Fox, J.C. (1986): Aggregate resources of the Bistcho Lake map area, NTS 84M/6-16; Alberta Resource Council, Map A84M05-16, scale 1:250 000.
- Paulen, R.C., Kowalchuk, C.J., Plouffe, A., Ward, B.C. and Smith, I.R. (2006): Surficial Geology of the Zama City Area, Alberta (NTS 84M/SE); Alberta Energy and Utilities Board, EUB/AGS Map 361 and Geological Survey of Canada, Open File 5184, scale 1:100 000.
- Plouffe, A., Paulen, R.C., Smith, I.R. (2006): Surficial geology, Thirteenth Creek, Alberta (NTS 84M/NW); Geological Survey of Canada, Open File 5070 and Alberta Energy and Utilities Board, EUB/AGS Map 395, scale 1:100 000.
- Smith, I.R., Paulen, R.C., Plouffe, A., Kowalchuk, C.J. and Peterson, R. (2005): Surficial mapping and aggregate resource assessment in northwest Alberta, in Summary of Activities 2005, British Columbia Ministry of Energy and Mines, p. 80-95.
- Smith, I.R., Plouffe, A. and Paulen, R.C. (2006): Surficial geology, Mega River, Alberta (NTS 84M/SW); Geological Survey of Canada, Open File 5237 and Alberta Energy and Utilities Board, EUB/AGS Map 396, scale 1:100 000.
- Zoltai, S.C. (1993): Cyclic development of permafrost in the peatlands of northwestern Alberta, Canada; Arctic and Alpine Research, v. 25, p. 240-246.

This is a common map legend for the surficial geology of northern Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

UNIT	UNIT NAME	DESCRIPTION AND GENESIS
QUATERNARY HOLOCENE		
A	ANTHROPOGENIC MATERIALS:	Culturally made or modified geological materials such that their physical properties (e.g., structure, cohesion, compaction) have been drastically altered.
O	ORGANIC DEPOSITS:	Undifferentiated peat (woody to fibrous muck) occurring in undifferentiated wetlands; commonly underlain by fine-grained, poorly drained glaciolacustrine deposits; includes marshes, swamps, bogs and fens.
OB	Bog peat:	Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is dominated by sphagnum mosses, heath shrubs and short, stunted trees.
OF	Fen peat:	Occurs in a peatland with water table at surface and slow internal drainage; peatland surface is dominated by sedges, with grasses and reeds near local pools, and is sparsely treed.
C	COLLUVIAL DEPOSITS:	Materials that have reached their present position as a result of direct, gravity-induced movement; commonly occurs as slope and slump deposits confined to valley slopes and floors; includes pre-existing bedrock, till, glaciolacustrine, glaciofluvial and eolian sediments, generally poorly sorted.
F	FLUVIAL DEPOSITS:	Sediments transported and deposited by streams and rivers; synonymous with alluvial. Includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g., postglacial floodplains, terraces, fans and deltas).
L	LACUSTRINE DEPOSITS:	Sediments deposited in and adjacent to recent and modern lakes; offshore sand, silt and clay, minor organic deposits; littoral (nearshore) beaches and bars; sand, silt and minor gravel.
E	EOLIAN DEPOSITS:	Wind-deposited sediments; well-sorted, medium to fine-grained sand and minor silt (loess); generally massive to locally cross bedded or ripple laminated; includes both active and vegetated deposits.
PLEISTOCENE		
LG	GLACIOLACUSTRINE DEPOSITS:	Primarily fine-grained, distal sediments deposited in or along the margins of glacial lakes, including sediments released by the melting of floating ice. Includes laminated (rhythmically bedded) to massive fine sand, silt and clay, and may contain ice-raftered stones.
LGL	Littoral and nearshore sediments:	Massive to stratified, well-sorted silty sand, pebbly sand and minor gravel; occurs as beaches, bars, spits and foreset deltaic deposits deposited during regression and lowering of glacial lakes.
FG	GLACIOFLUVIAL DEPOSITS:	Sediments deposited by glacial meltwater streams as subaerial or subaqueous outwash. Includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Features include meltwater channels, kettle holes, terraces and minor ice-contact sediments.
FGI	Ice-contact sediments:	Sediments deposited by glacial meltwater streams in direct contact with glacial ice, either in front of (kame terraces) or within (eskers, crevasse ridges) glacial ice. Includes massive to stratified, poor to moderately sorted, coarse sediments (predominantly pebble gravel and coarse sand, locally till) and may show evidence of ice melting (slumped structures).
M	MORAINE:	Nonsorted diamicton (till) deposited directly by glacial ice consisting of a mixture of clay, silt, sand and minor pebbles, cobbles and boulders. Locally, this unit may contain blocks of bedrock, pre-existing stratified sediment and till, and lenses of glaciolacustrine and/or glaciofluvial sediment.
MS	Stagnant ice moraine:	Material resulting from the collapse and lateral movement of englacial and supraglacial sediment in response to melting (ablation) of buried stagnant ice at the ice margin; sediment is mainly diamicton, but locally includes stratified sediments of glaciolacustrine or glaciofluvial origin. Characterized by low to high-relief hummocky topography.
MT	Ice-thrust moraine:	Terrain resulting from glacio-tectonic transport of originally subglacial material deposited by the glacier more or less intact; deposits may include syngenetic till, as well as masses of pre-existing till, stratified drift and/or bedrock. Characterized by high to moderate relief and features include hill-hole pairs and glacio-tectonic moraine ridges.
MF	Fluted moraine:	Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smoothed hills; all landforms parallel the local ice flow direction; includes flutes, drumlins and drumlinoids.
FP	PREGLACIAL FLUVIAL DEPOSITS:	Sediments transported and deposited by streams and rivers prior to glaciation. Includes sand and gravel deposits occurring in paleovalleys (i.e., preglacial floodplains, terraces, fans and deltas); ranging from middle Wisconsin to late Tertiary.
PRE-QUATERNARY		
RT	UNCONSOLIDATED FLUVIAL GRAVELS:	Predominately well-sorted, quartzite and chert gravel and cobbles; Cordilleran source, Paleogene (Tertiary) to early Quaternary.
R	BEDROCK:	Undifferentiated; may include clastic sedimentary rock, shale, coal, carbonate and crystalline (Shield), kimberlite and/or coal.

UNIT NOTATION

Example: GLACIOLACUSTRINE plain



Textural Modifier

Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture; i.e., sc for sandy clay

p = pebble

g = gravel

s = sand

\$ = silt

c = clay

a = sand-silt-clay

GENETIC & GEOMORPHIC MODIFIERS

- | | | |
|---|---------------------------|---|
| c | crevasse fill | ice-contact ridges, ice-squeeze deposits and linear forms deposited by meltwater in stagnant ice |
| d | doughnut rings and ridges | circular hummocks with a central depression, plateau mounds and brain-like pattern ridges, low to moderate relief |
| e | eroded | planar surface eroded by glacial meltwater, often capped by a boulder lag deposit and/or thin deposit of sand and gravel |
| f | fan | gently sloping fan-shaped mass of detrital debris |
| g | gullied | slopes dissected by modern ravines created by intermittent runoff |
| h | hummock | assemblage of approximately equidimensional hills and hollows; moderate to high relief (commonly greater than 2 m) |
| k | collapse | depression, including kettles, pitted morphology, thermokarst depressions, karst sinkholes |
| m | meander | sinuous curves, loops and oxbows produced as meltwater and modern streams shift their channels over time |
| p | plain | deposit greater than 2 m thick; commonly masks geomorphic pattern of underlying deposits; flat to gently rolling topography (commonly less than 2 m relief) |
| r | ridged | one or more parallel or subparallel, convex, linear morphological elements with a length-to-width ratio greater than 2; low to high relief |
| s | slumped | landslide blocks, slope failure debris |
| t | terrace | terrace bench cut by either meltwater or wave action; antiplanation terrace, kame terrace |
| u | undulating | low-relief rolling terrain; swell and swale topography |
| v | veneer | thin mantle of unconsolidated material too thin to mask the minor irregularities of the surface of the underlying material; it ranges in thickness from 10 cm to 1 metre and may be discontinuous |
| w | washboard | low-relief transverse moraine ridges, usually formed from basal ice shearing |
| y | dissected | channelled or dissected by glacial meltwater flow; dissected terrain by Holocene fluvial activity |
| z | delta | lake delta; ice-contact delta |

Complex

Where two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the combination is given in a two or three position designation set off by slashes denoting arbitrary percentage limits. For example,

'Mp/LGv' means the area is underlain by approximately 60% morainal plain and up to 40% glaciolacustrine veneer.

'Mv/LGv/FGp' means at least 60% of the area is underlain by morainal veneer, with up to 40% glaciolacustrine veneer and less than 15% glaciofluvial plain.

'LGp//M' means more than 60% of the area is underlain by a glaciolacustrine plain, with less than 15% moraine.

Stratigraphic Sequence

Where materials of different origins or textures are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:

'sLGv | Mp' Thin sandy glaciolacustrine sediment deposited on morainal plain

Transitional Association

Locally, two or more terrain units are juxtaposed by reason of related origin, temporal sequence or ambiguous geomorphic distinction. In the last case, both components may be present. Such situations are identified by a compound designation marked by a hyphen. Examples: 'FGz-LGz' indicating ice-contact delta indistinguishable from glaciolacustrine delta, or 'FGlk-MSh' indicating ice-contact kame and kettle topography that blends with hummocky stagnant ice moraine.

Morphologic Overprint

Where a sequence of geomorphic processes has produced a multi-aspect or compound terrain fabric, the geomorphic modifier suffixes are appended in the inferred order of superposition. 'Mpry' means a plain of till has been moulded into ridge forms and finally dissected by modern streams. 'FGphr' means a glaciofluvial plain has been discontinuously covered by ice-contact hummocks and ridges.