

This is a common map legend for the surficial geology of northern Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

UNIT	UNIT NAME	DESCRIPTION AND GENESIS
<b>QUATERNARY HOLOCENE</b>		
A	ANTHROPOGENIC MATERIALS:	Artificially made ground or geological materials that have been disturbed by human activity, such that their physical properties (e.g. structure, cohesion, compaction) have been drastically altered.
O	ORGANIC DEPOSITS:	Undifferentiated peat (woody to fibrous muck) occurring in wetlands; commonly underlain by fine-grained, poorly drained glaciolacustrine deposits; includes marshes, swamps, bogs and fens.
OB	Bog peat:	Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is dominated by sphagnum mosses, heath shrubs and short, stunted trees.
OF	Fen peat:	Occurs in peatland which receives water from slowly flowing streams and groundwater, with the water table lying at the land surface; peatland surface is dominated by sedges, with grasses and reeds near local pools, and is sparsely treed.
C	COLLUVIAL DEPOSITS:	Materials that have reached their present position as a result of direct, gravity-induced movement; commonly occurs as slope and slump deposits confined to valley slopes and floors; includes pre-existing bedrock, till, glaciolacustrine, glaciofluvial and eolian sediments, generally poorly sorted.
F	FLUVIAL DEPOSITS:	Sediments transported and deposited by streams and rivers; synonymous with alluvium. Includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g., postglacial floodplains, terraces, fans and deltas).
L	LACUSTRINE DEPOSITS:	Sediments deposited in and adjacent to recent and modern lakes. Includes offshore sand, silt and clay; minor organic deposits; may also include minor littoral (nearshore) beaches and bars composed of sand, silt and minor gravel.
E	EOLIAN DEPOSITS:	Wind deposited sediments; comprise silt, medium- to fine-grained sand and minor silt; generally massive to locally cross-bedded or ripple-laminated; includes both active and vegetated dunes and sand sheets.
<b>PLEISTOCENE</b>		
LG	GLACIOLACUSTRINE DEPOSITS:	Primarily fine-grained, distal sediments deposited in or along the margins of glacial lakes, including sediments released by the melting of floating ice. Includes laminated (rhythmically bedded) to massive fine sand, silt and clay, and may contain ice-rafted debris.
LGL	Littoral and nearshore sediments:	Massive to stratified, well-sorted silty sand, pebbly sand and minor gravel; occurs in beaches, bars, spits and deltaic features deposited during regression and lowering of glacial lakes.
FG	GLACIOFLUVIAL DEPOSITS:	Sediments deposited by glacial meltwater streams as subaerial or subaqueous outwash. Includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Features include meltwater channels, kettle holes, terraces and minor ice-contact sediments.
FGI	Ice-contact sediments:	Sediments deposited by meltwater streams flowing either in direct contact with the ice margin (kame terraces) or within and/or under glacial ice (eskers, crevasse ridges). Includes massive to stratified, poor to moderately sorted, coarse-grained sediments (predominantly pebble gravel and coarse-grained sand, locally silt) and may show evidence of ice melting (slumped structures).
M	MORAINES:	Diamictic (til) deposited directly by glacial ice and consisting of a mixture of clay, silt, sand and minor pebbles, cobbles and boulders. Locally, this unit may contain blocks of bedrock, pre-existing stratified sediment and till, or lenses of glaciolacustrine and/or glaciofluvial sediment.
MS	Stagnant ice moraine:	Material resulting from the collapse and slumping of englacial and supraglacial sediment in response to the melting of stagnant ice at the ice margin; sediment is mainly diamictic, but locally includes stratified sediments of glaciolacustrine or glaciofluvial origin. Characterized by low to high-relief hummocky topography.
MT	Ice-thrust moraine:	Terrain formed from the glacio-tectonic displacement of materials as blocks or rafts in a more or less intact state. Materials may include syngenetic till, as well as masses of pre-existing sediments and/or bedrock. Characterized by high to moderate relief and features include hill-hole pairs and glacio-tectonic moraines.
MF	Fluted moraine:	Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smooth hills; all landforms parallel the local ice flow direction; includes flutes, drumlins and drumlinoids.
FP	PREGLACIAL FLUVIAL DEPOSITS:	Sediments transported and deposited by streams and rivers prior to glaciation. This includes sand and gravel deposited in paleovalleys (i.e., preglacial floodplains, terraces, fans and deltas).
<b>PRE-QUATERNARY</b>		
RT	UNCONSOLIDATED FLUVIAL GRAVELS:	Predominately well-sorted, quartzite and chert gravel and cobbles; Cordilleran source, Paleogene to Neogene.
R	BEDROCK	

SYMBOL LEGEND		BASE MAP LEGEND	
Landslide and active layer failure scar (small)	↓	Road-paved-primary	—
Landslide and active layer failure scar (large)	—	Road-gravel-primary	—
Eolian forms; dune ridges	—	Road-unimproved	—
Beach or strandline	—	Trail-truck	—
Wave-cut bench	—	River	—
Escarpment	—	Lake	—
Melwater channel (minor)	—	UTM, Zone 11 Grid	+
Melwater channel (minor, flow indicated)	—	Contour, intervals 50 metres	—
Melwater channel (major)	—	Town	—
Melwater channel (major)	—		
Crevasse filling	—		
Kettle	—		
Esker (paleoflow direction unknown)	—		
Drumlinoid or streamlined landform	—		
Minor moraine ridge	—		
Major moraine ridge	—		
Ice thrust ridge	—		
Bedrock outcrop	X		
Gravel and/or sand pit	—		
Section of stratigraphic interest	—		

**UNIT NOTATION**  
Example: sandy GLACIOLACUSTRINE plain

**Textural Modifier**  
Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture; i.e., sc for sandy clay.

**GENETIC & GEOMORPHOLOGICAL MODIFIERS**

**Complex**  
If two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the combination is given in a two- or three-position designation set off by slashes denoting arbitrary percentage limits. Examples are:

**Stratigraphic Sequence**  
If materials of different origins or textures are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:

**Transitional Association**  
Locally, two or more terrain units are juxtaposed by reason of related origin, temporal sequence or ambiguous geomorphological distinction. In the last case, both components may or may not be present. Such situations are identified by a compound designation marked by a hyphen. Examples are:

**Morphological Overprint**  
If a sequence of geomorphological processes has produced a multi-aspect or compound terrain fabric, the geomorphological modifier suffixes are appended in the inferred order of superposition. Mpry indicates a moraine plain has been moulded into ridges and finally dissected by streams. FGpr indicates a glaciofluvial plain that includes discontinuous hummocks and ridges.


**Methodology**  
The Alberta Geological Survey completed the surficial mapping in 2010. Observations made during field mapping were combined with the interpretation of Light Detection and Ranging (LIDAR) data (Figure 1), supplied by Alberta Sustainable Resource Development, and digital orthorectified airphotos (1:50 000 scale). The LIDAR digital elevation model (DEM) was used to delineate landforms through shaded-relief images created from three illumination directions. The shaded relief shown on the map was produced by using shaded-relief (315° illumination azimuth, 45° inclination) and slope-gradient images.

**Acknowledgements**  
G. Ouellette, N. Atkinson, J.G. Pawlowicz, G.J. Prior, J.A. Weiss, D.C. Hay and K. McKay assisted with fieldwork. K. McKay completed the digital cartography and GIS. Spatial Data Warehouse Ltd. provided the base data. N. Atkinson provided constructive comments that improved this map.

**References**  
Atkinson, N. (2008): Surficial geology of the High Prairie area (NTS 83N/SE), Energy Resources Conservation Board, ERC/BAGS Map 421, scale 1:100 000.  
Mathews, W.H. (1980): Retreat of the last ice sheets in northeastern British Columbia and adjacent Alberta; Geological Survey of Canada, Bulletin 331, 22 p.  
Paulen, R.C., Pawlowicz, J.G. and Fenton, M.M. (2004): Surficial geology of the Willow River area (NTS 830/NE), Alberta Energy and Utilities Board, EUB/AGS Map 313, scale 1:100 000.  
Atkinson, N. (2010): Surficial geology of the Faust area (NTS 830/SW), Energy Resources Conservation Board, ERC/BAGS Map 554, scale 1:100 000.  
St-Onge, D.A. (1972): Sequence of glacial lakes in north-central Alberta; Geological Survey of Canada, Bulletin 213, 142 p.  
St-Onge, D.A. (1975): Surficial geology, west of fifth meridian, Whitecourt, Alberta; Geological Survey of Canada, Map 1367A, scale 1:250 000.

Alberta Geological Survey  
(780) 422-1927  
www.ags.gov.ab.ca  
Published 2010  
ISBN 978-0-7785-9639-5

**Map 553**  
**Surficial Geology of the Lesser Slave River Area (NTS 830/SE)**  
Geology by: S.M. Pawley



Scale 1:100 000  
Projection: Universal Transverse Mercator  
Datum: North American Datum, 1983

