

This is a common map legend for the surficial geology of northern Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

UNIT	UNIT NAME	DESCRIPTION AND GENESIS
QUATERNARY HOLOCENE		
A	ANTHROPOGENIC MATERIALS:	Artificially made ground or geological materials that have been disturbed by human activity, such that their physical properties (e.g. structure, cohesion, compaction) have been drastically altered.
O	ORGANIC DEPOSITS:	Undifferentiated peat (woody to fibrous muck) occurring in wetlands; commonly underlain by fine-grained, poorly drained glaciolacustrine deposits; includes marshes, swamps, bogs and fens.
OB	Bog peat:	Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is dominated by sphagnum mosses, heath shrubs and short, stunted trees.
OF	Fen peat:	Occurs in peatland which receives water from slowly flowing streams and groundwater, with the water table lying at the land surface; peatland surface is dominated by sedges, with grasses and reeds near local pools, and is sparsely treed.
C	COLLUVIAL DEPOSITS:	Materials that have reached their present position as a result of direct, gravity-induced movement; commonly occurs as slope and slump deposits confined to valley slopes and floors; includes pre-existing bedrock, till, glaciolacustrine, glaciolacustrine and eolian sediments, generally poorly sorted.
F	FLUVIAL DEPOSITS:	Sediments transported and deposited by streams and rivers; synonymous with alluvium. Includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g., postglacial floodplains, terraces, fans and deltas).
L	LACUSTRINE DEPOSITS:	Sediments deposited in and adjacent to recent and modern lakes; includes offshore sand, silt and clay, minor organic deposits; may also include minor littoral (nearshore) beaches and bars composed of sand, silt and minor gravel.
E	EOLIAN DEPOSITS:	Wind deposited sediments; comprise well-sorted, medium- to fine-grained sand and minor silt; generally massive to locally cross-bedded or ripple-laminated; includes both active and vegetated dunes and sand sheets.
PLEISTOCENE		
LG	GLACIOLACUSTRINE DEPOSITS:	Primarily fine-grained, distal sediments deposited in or along the margins of glacial lakes, including sediments released by the melting of floating ice. Includes laminated (rhythmically bedded) to massive fine sand, silt and clay, and may contain ice-rafted debris.
LGL	Littoral and nearshore sediments:	Massive to stratified, well-sorted silty sand, pebbly sand and minor gravel; occurs in beaches, bars, spits and deltaic forests deposited during regression and lowering of glacial lakes.
FG	GLACIOFLUVIAL DEPOSITS:	Sediments deposited by glacial meltwater streams as subaqueous or outwash. Includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Features include meltwater channels, kettle holes, terraces and minor ice-contact sediments.
FGI	Ice-contact sediments:	Sediments deposited by meltwater streams flowing either in direct contact with the ice margin (kame terraces) or within and/or under glacial ice (eskers, crevasse ridges). Includes massive to stratified, poorly to moderately sorted, coarse-grained sediments (predominantly pebbly gravel and coarse-grained sand, locally till) and may show evidence of ice melting (slumped structures).
M	MORAINES:	Diamicton (till) deposited directly by glacial ice and consisting of a mixture of clay, silt, sand and minor pebbles, cobbles and boulders. Locally, the unit may contain blocks of bedrock, pre-existing stratified sediment and till, or lenses of glaciolacustrine and/or glaciolacustrine sediment.
MS	Stagnant ice moraine:	Material resulting from the collapse and slumping of englacial and supraglacial sediment in response to the melting of buried stagnant ice at the ice margin; sediment is mainly diamict, but locally includes stratified sediments of glaciolacustrine or glaciolacustrine origin. Characterized by low to high-relief hummocky topography.
MT	Ice-thrust moraine:	Terrain formed from the glaciotectionic displacement of materials as blocks or rafts in a more or less intact state. Materials may include syngenetic till, as well as masses of pre-existing sediments and/or bedrock. Characterized by high to moderate relief and features include hill-hole pairs and glaciotectionic moraines.
MF	Fluted moraine:	Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smoothed hills; all landforms parallel the local ice flow direction; includes flutes, drumlins and drumlinoids.
FP	PREGLACIAL FLUVIAL DEPOSITS:	Sediments transported and deposited by streams and rivers prior to glaciation. This includes sand and gravel deposited in paleovalleys (i.e., preglacial floodplains, terraces, fans and deltas).
PRE-QUATERNARY		
RT	UNCONSOLIDATED FLUVIAL GRAVELS:	Predominantly well-sorted, quartzite and chert gravel and cobbles; Cretaceous source, Paleogene to Neogene.
R	BEDROCK	

UNIT NOTATION
Example: sandy GLACIOLACUSTRINE plain

Textural modifier	Genetic unit	Geomorphic modifier
s	LG	p

Textural Modifier
Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture. I.e., sc for sandy clay.
p = pebble
g = gravel
s = sand
\$ = silt
c = clay
a = sand-silt-clay

GENETIC & GEOMORPHOLOGICAL MODIFIERS

c	crevasse fill	ice-contact ridges formed by the slumping of sediment into crevasses on the ice surface or the squeezing of till into fractures at the ice base
d	doughnut rings and ridges	circular hummocks with a central depression, plateau mounds and brain-like pattern ridges; low to moderate relief
e	eroded	planar surface eroded by glacial meltwater, often capped by a boulder lag and/or thin deposit of sand and gravel
f	fan	gently sloping fan-shaped mass of detrital debris
g	gullied	slopes dissected by modern ravines created by intermittent runoff
h	hummock	assemblage of approximately equidimensional hills and hollows; moderate to high relief (commonly greater than 2 m)
k	kollapse	depression, includes kettle holes, pitted morphology, thermokarst depressions, karst sinkholes
kt	thermokarst	topography consists of mounds and circular, poorly drained lowlands; formed by the local melting of ground ice and subsequent settling of the land surface; often associated with ice-rich permafrost occurring in thick peat blankets that overlie fine-grained moraine
m	meander	sinuous curves, loops and oxbows produced as meltwater and modern streams shift their channels over time
p	plain	deposit greater than 2 m thick; commonly masks geomorphic pattern of underlying deposits; flat to gently rolling topography (commonly less than 2 m relief)
r	ridged	one or more parallel or subparallel, convex, linear morphological elements with a length-to-width ratio greater than 2; low to high relief
s	slumped	landslide blocks, slope failure debris
t	terrace	a bench of either erosional or depositional origin that flanks the sides of floodplains, valleys and lakes; includes fluvial and glaciolacustrine terraces, shoreline terraces and topographic terraces
u	undulating	low-relief rolling terrain; swell and swale topography
v	vener	thin mantle of unconsolidated sediment that is too thin to mask the minor irregularities of the surface of the underlying material; it ranges in thickness from 10 cm to 1 metre and may be discontinuous
y	dissected	channelled or dissected by glacial meltwater and/or Holocene fluvial activity
z	delta	lake delta; ice-contact delta

Complex
Where two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the combination set off by a slash between arbitrary percentage limits. Examples are:

'Mp/LGv'	indicates the area is underlain by approximately 60% morainal plain and up to 40% glaciolacustrine veneer
'Mv/LGvGp'	indicates at least 60% of the area is underlain by morainal veneer, with up to 40% glaciolacustrine veneer and less than 15% glaciolacustrine plain
'LGP/IM'	indicates more than 60% of the area is underlain by a glaciolacustrine plain, with less than 15% moraine

Stratigraphic Sequence
Where materials of different origins or textures are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:

'sLGV M'	indicates sandy glaciolacustrine veneer deposited on morainal plain
------------	---

Transitional Association
Locally, two or more terrain units are juxtaposed by reason of related origin, temporal sequence or ambiguous geomorphological distinction. In the last case, both components may or may not be present. Such situations are identified by a compound designation marked by a hyphen. Examples are:

'FGz-LGz'	indicates ice-contact delta indistinguishable from glaciolacustrine delta
'LG-LGL'	indicates glaciolacustrine indistinguishable from littoral and nearshore glaciolacustrine sediment

Morphological Overprint
Where a sequence of geomorphological processes has produced a multi-aspect or compound terrain fabric, the geomorphological modifier suffixes are appended in the inferred order of superposition. 'Mpy' indicates a morainal plain that has been moulded into ridges and finally dissected by streams. 'FGpr' indicates a glaciolacustrine plain that includes discontinuous hummocks and ridges.

Methodology
Alberta Geological Survey conducted surficial geology field mapping in the area in 2006, 2007, and 2010. Observations made during field mapping were combined with the interpretation of Light Detection and Ranging (LiDAR) bare-earth data (Figure 1), SPOT-5 panchromatic imagery, and image classification of peatlands from Landsat 8 multispectral data (courtesy of the U.S. Geological Survey). Landforms were initially delineated using an object-based, image segmentation approach, performed on a slope derivative calculated from the LiDAR Digital Elevation Model (DEM). These landforms were classified into several categories using terrain derivatives calculated from the LiDAR DEM (e.g., slope, ruggedness, texture). This initial classification was corrected and augmented by further interpretation and digitization using shaded-relief images created from three illumination directions. The shaded relief shown as an underlay on the main map was produced by fusing shaded-relief (315° illumination azimuth, 45° declination) and slope-gradient images.


Acknowledgements
Fieldwork during the 2006-2007 season was performed by R.C. Paulen, assisted by S. Bottenli and K. McKay. N. Atkinson assisted with the fieldwork during 2010. K. McKay completed the digital cartography and GIS. Spatial Data Warehouse Ltd. provided the base data. Environment and Sustainable Resource Development provided the DEM. D.J. Utting provided constructive comments that improved this map.

References
Fenton, M.M., Waters, E.J., Pawley, S.M., Atkinson, N., Utting, D.J. and McKay, K. (2013). Surficial geology of Alberta; Alberta Energy Regulator, AER/AGS Map 601, scale 1:1 000 000.
Pawley, S.M. (2013). Surficial Geology of the Grand Rapids Area (NTS 84A/SE); Alberta Energy Regulator, AER/AGS Map 566, scale 1:100 000.
Pawley, S.M. (2013). Surficial Geology of the Wabasca Area (NTS 84A/SW); Energy Resources Conservation Board, ERCB/AGS Map 561, scale 1:100 000.
Pawley, S.M. (2011). Surficial Geology of the Thickwood Hills Area (NTS 84A/NE); Energy Resources Conservation Board, ERCB/AGS Map 555, scale 1:100 000.

Fisher, T.G., Waterson, N., Lowell, T.V. and Hajdas, I. (2009). Deglaciation ages and meltwater routing in the Fort McMurray region, northeastern Alberta and northwestern Saskatchewan, Canada; Quaternary Science Reviews, 28, p. 1609-1624.

Recommended Reference Format
Pawley, S.M. (2014). Surficial Geology of the Chipewyan Lake Area (NTS 84A/NW); Alberta Energy Regulator, AER/AGS Map 572, scale 1:100 000.


SYMBOL LEGEND	BASEMAP LEGEND
Permafrost; relict and/or active	Primary road, gravel
Eolian forms; dune ridges	Unimproved road
Beach or strandline	Truck trail
Wave-out bench	River
Meltwater channel (minor)	Lake
Meltwater channel (major)	UTM, Zone 12 Grid
Crevasse filling	Contour, intervals 50 metres
Ice thrust ridge	Town
Esker (paleoflow direction unknown)	
Drumlinoid or streamlined landform	
Buried drumlinoid or streamlined landform	
Minor moraine ridge	
Major moraine ridge	



Alberta Geological Survey
(780) 422-1927
www.ag.srv.ab.ca
Published 2014
ISBN 978-1-4601-0127-8

Map 572
Surficial Geology of the Chipewyan Lake Area (NTS 84A/NW)

Geology by: S.M. Pawley



Projection: Universal Transverse Mercator
Datum: North American Datum, 1983

